

## Mayo Clinic Urology postoperative opioid prescribing guidelines for opioid-naïve patients

Procedures with similar levels of perceived pain (based on a critical assessment of historical data) were grouped into four tiers with an agreed-upon maximum recommended quantity of opioids.

<b>Proposed level</b>	<b>Representative procedures</b>	<b>Historical median oral morphine equivalent (OME) prescribed</b>	<b>Proposed max OME</b>	<b>Tramadol (50 mg)</b>	<b>Hydrocodone (5 mg)</b>	<b>Oxycodone (5 mg)</b>	<b>Hydromorphone (2 mg)</b>
<b>0</b>	Vasectomy	0	0	-	-	-	-
<b>1.0</b>	Circumcision (adult)	150	50-75	10-15	10-15	7.5-12	7.5-12
	Transurethral resection of bladder tumor	150	50-75				
	Holmium laser enucleation of prostate	113	50-75				
	Photovaporization of the prostate	100	50-75				
	Cystolitholapaxy	100	50-75				
	Ureteral stent placement	100	50-75				
<b>1.1</b>	Ureteroscopic stone extraction	150	100	20	20	15	15
<b>2</b>	Robotic-assisted laparoscopic partial nephrectomy	225	200	40	40	25	25
	Hand-assisted laparoscopic nephrectomy	225	200				
	Robotic-assisted laparoscopic radical prostatectomy	225	200				
	Artificial genitourinary sphincter	225	200				
	Hydrocelectomy/Spermatocelectomy	200	200				
	Percutaneous nephrolithotomy	225	200				
	Inflatable penile prosthesis/malleable penile prosthesis placement	225	200				
	Autologous pubovaginal sling	200	200				
<b>3</b>	Open radical retropubic prostatectomy	300	300	60	60	40	40
	Open radical nephrectomy	300	300				
	Open partial nephrectomy	240	300				
	Open cystoprostatectomy with ileal conduit/ileal neobladder	300	300				
	Retroperitoneal lymph node dissection (LND)	300	300				